

Why Bigger Areas Hold More Species

Decoding the Species-Area Relationship:
From Observation to Ecological Law




Alexander von Humboldt

During his pioneering explorations of South American jungles in the 19th century, this great German naturalist and geographer noticed a consistent spatial pattern in biodiversity.

Within a region, species richness increases with increasing explored area, but only up to a limit.

The Intuitive Rule of Diminishing Returns

As we survey larger tracts of land, we discover new species rapidly at first. However, the rate of discovery eventually slows down.





You've Learned the Theory. Let's Test Them Now.

$$E=mc^2$$



Turn theory into marks with **unlimited MCQs, smart analysis, and focused revision.**

Practice Now




At prep.topall.app

Key Highlights

- » Free Study Material, Videos & Infographics
- » Push Your Limits with Level-Based Challenges
- » Climb the Leaderboard Among Fellow Aspirants
- » Practice Unlimited MCQs for NEET & JEE
- » Get Clear Explanations, Not Just Marks
- » Track Your Performance Smarter




The Raw Shape: Rectangular Hyperbola



Why does it curve?

The flattening represents the biological limit of new species discovery. Eventually, you run out of unique habitats and unrecorded species within a bounded region.

Straightening the Curve for Analysis



By plotting the data on a logarithmic scale, the curving relationship becomes a straight line. This mathematical transformation allows ecologists to instantly compare the scaling rules of completely different ecosystems just by looking at the slope.

The Mathematical Engine

S = Species Richness

The total number of species found

Z = Slope of the line

The regression coefficient, defining how fast species accumulate

$$\log S = \log C + Z \log A$$

C = Y-intercept

The baseline species count for a unit area

A = Area

The geographical size of the surveyed region

The Z-Value: Scaling Across Worlds

Regional Scale

0.1 to 0.2



Remains amazingly similar regardless of the taxonomic group or the region.


Plants in Britain

Birds in California

Molluscs in New York

Continental Scale


0.6 to 1.2



Slopes become much steeper when analyzing vast, continent-wide areas.

Frugivorous birds & mammals in tropical forests ($Z = 1.15$)

Figure 13.2: Species-Area Relationship



This dual-view model demonstrates how the natural, physical bounds of biodiversity (the hyperbola) translate into a predictable, comparative mathematical rule (the straight line on a log scale).



You've Learned the Theory.

Now Let's Test It.

Reading notes gets you knowledge.

Practicing MCQs gets you marks.

That's exactly what TopAll Pro is built for.

Try TopAll Pro Now

What TopAll Pro Does Differently

-  **Stuck? Snap it. Solve it. — Doubts AI**
Photo → step-by-step solution in seconds, 24/7
-  **Your AI Coach finds your blind spots**
Concept-level weakness detection — not just chapter-level
-  **4.5 Lakh+ questions. Never run out.**
13× more content than any other prep app in India
-  **Build your own test in 60 seconds**
Pick a chapter, set difficulty — AI does the rest instantly.
-  **AI Recommendation that helps you improve**
AI Recommendation Analysis your Tests and tell you where you need to improve.



Practice the moment you finish studying
Jump in while the concept is still fresh in your head

4.5L+
MCQs

1.5L+
Students

AI Powered
In-Depth Analysis